

عنوان مقاله:

Perineal Leiomyoma: A Case Report

محل انتشار:

فصلنامه زنان و مامایی و سرطانهای زنان، دوره 5، شماره 2 (سال: 1399)

تعداد صفحات اصل مقاله: 3

نویسندگان:

Sarmad Nourooz Zadeh - *Medical Doctor, School of Medicine, Urmia University of Medical Sciences, Urmia, Iran*

Sedigheh Ghasemian Dizajmehr - *Department of Obstetrics and Gynecology, School of Medicine, Urmia University of Medical Sciences, Urmia, Iran*

Farzaneh Rashidi Fakari - *PhD, Student Research Committee, Department of Midwifery and Reproductive Health, School of Nursing and Midwifery, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

Hania Fattahi - *Resident of Pathology, School of Medicine, Urmia University of Medical Sciences, Urmia, Iran*

Mohsen Ghasemian - *Medical Doctor, School of Medicine, Ardabil University of Medical Sciences, Ardabil, Iran*

خلاصه مقاله:

Introduction: Leiomyomas are tumors of the soft tissues. The incidence of myomas within the perineum is absolutely rare and only a few reports have been made on the matter. We herein report a case of perineal myoma in a virgin woman, which was successfully excised. Case report: We present a 35-year-old virgin woman, with complaint referred to a mass in her perineum which had first exhibited signs 5 years prior to the time she referred to us and had increased in size in a steady manner ever since. The mass had caused no gynecologic, rectal, or urinary symptoms. Clinical examination revealed a painless, mobile mass (6cm*6cm) with [unknown consistency] in the right perineum with extension to the distal of the labium majus of the same side. Conclusion: Treatment of symptomatic leiomyomas relies on surgical excision of the mass. However, the surgical method of choice is a matter of debate in previous studies.

کلمات کلیدی:

Leiomyoma, Perineum, Tissue

لینک ثابت مقاله در پایگاه سیویلیکا:

<https://civilica.com/doc/1138642>

